

## Acetaminophen Dosage (Tylenol®)

**If your infant is under 2 months of age, call the office immediately for fever greater than 100.4°F  
 (617) 471-3411**

A Dose is by weight given every 4 hours		Infants' Drops 160mg/5mL	Children's Suspension Liquid 160mg/5mL	Children's Soft Chews-Chewable Tablets 80mg each	Junior Strength Chewable Tablets 160mg each	Feverall (Suppositories) 160mg/5mL
Weight	Age	Dropperful <i>(Use dropper)</i>	Teaspoon <i>(Use only the dosing cup provided)</i>	Tablet	Tablet	Suppository
6-11 lbs.	0-3 Months	1.25mL				
12-17 lbs.	4-11 Months	2.5mL	½ tsp.			¾ (120mgm)
18-23 lbs.	12-23 Months	3.75mL	¾ tsp.			1 (120mgm)
24-35 lbs.	2-3 Years	5mL	1 tsp.	2		1 ½ (120mgm)
36-47 lbs.	4-5 Years		1 ½ tsp.	3		2 (120mgm)
48-59 lbs.	6-8 Years		2 tsp.	4	2	1 (325mgm)
60-71 lbs.	9-10 Years		2 ½ tsp.	5	2 ½	1 ¼ (325mgm)
72-95 lbs.	11 Years		3 tsp.	6	3	1 ½ (325mgm)
96 lbs. & Over	12 Years				4	2 (325mgm)

**Please note: One does last 4 hours. Do not give more than 5 doses in any 24 hours.**

## Ibuprofen Dosage (Motrin® or Advil®)

A Dose is by weight given every 6-8 hours Do not give Motrin under 6 months of age		Infants' Ibuprofen Concentrated Drops 50mg/1.25mL	Children's Ibuprofen Suspension 100mg/5mL	Children's Soft Chewable Tablets 50mg each	Junior Strength Chewable Tablets 100mg each	Junior Strength Ibuprofen Caplets 100mg each
Weight	Age	Dropperful (Use dropper)	Teaspoon (Use only the dosing cup provided)	Tablet	Tablet	Caplet
	Under 6 Months					
12-17 lbs.	6-11 Months	1 = (1.25mL)				
18-23 lbs.	12-23 Months	1 ½ = (1.875mL)				
24-35 lbs.	2-3 Years		1 tsp.	2		
36-47 lbs.	4-5 Years		1 ½ tsp.	3		
48-59 lbs.	6-8 Years		2 tsp.	4	2	2
60-71 lbs.	9-10 Years		2 ½ tsp.	5	2 ½	2 ½
72-95 lbs.	11 Years		3 tsp.	6	3	3
96 lbs. & Over	12 Years		4 tsp.	8	4	4 Junior Caplets or 2 Adult Caplets

- Milliliter is abbreviated as mL; 5mL equals one teaspoon (tsp.)
- Don't use household teaspoons, which can vary in size.
- Aspirin should not be used in children to treat fever or pain.